

Supplementary Files

[Iranian Journal of Weed Science: 2024; 20\(1\): 115-131](#)

The Response of Some of Alien Weeds in Iran to Environmental Factors and Their Distribution: A Literature Review

Sima Sohrabi^{1*}, Javid Gherekhloo², Mehdi Minbashi Moeini³, Eskandar Zand³

1. Faculty of Agriculture, Ferdowsi University of Mashhad, Mashhad, Iran.

2. Department of Plant Production, Gorgan University of Agricultural Sciences and Natural Resources, Gorgan, Iran.

3. Iranian Research Institute of Plant Protection, Agricultural Research, Education and Extension Organization (AREEO), Tehran, Iran.

Species	life	base	optimum	max	NaCl	Osmotic	PH	depth-p	INS
<i>Cuscuta campestris</i>	Annual	5.35	26.11	46.78	11.9	0.52	4.5 - 7.5	3	Naturalized
<i>Ipomoea hederacea</i> Jacq.	Annual	11.2	26.7	40	26	1.03	4-7.5	10	Invasive
<i>Ipomoea purpurea</i> L.	Annual	7.34	25.55	39.52	33.22	0.4	6-7.5	10	Naturalized
<i>Ipomoea lacunosa</i> L.	Annual	9.5	30.3	45.2	16.25	0.83	6 to 7.5	5	Naturalized
<i>Amaranthus retroflexus</i>	Annual	6.65	30.56	51.06	10.4	0.37	5 to 7	4	Naturalized
<i>Amaranthus hybridus</i>	Annual	14.37	34.21	44.5	35	0.64	4 to 6	2	Naturalized
<i>Amaranthus blitoides</i>	Annual	14.24	31.65	44.2	45	0.5	4.5 to 8.5	3	Naturalized
<i>Euphorbia maculata</i>	Annual	9.8	28	43.4	11.2	0.53	4 to 10	2	Invasive
<i>Euphorbia heterophylla</i>	Annual	11	34	45	4.5	0.67	5 to 7.5	4	Invasive
<i>Physalis angulata</i>	Annual	8.03	35.8	42.3	5.925	0.29	6 to 8	6	Naturalized
<i>Abutilon theophrasti</i> Medik.	Annual	5	34.84	48.16	11	0.3	4 to 9	6	Naturalized
<i>Datura innoxia</i> Mill.	perennial	6	25	40	6	0.27	5.5 to 6	2	Naturalized
<i>Ambrosia psilostachya</i> DC.	Perennial	5	25.5	42.13	10.4	0.24	5.7 to 7.9	10	Invasive
<i>Echinochloa oryzicola</i> (Vasinger) Vasinger	Annual	11	30	35	11.8	0.5	4.7-8.3	1.5	Naturalized

References

- (2015). 'Survey of Germination and Phenology of Dodder (*Cuscuta campestris*) in Sugar Beet (*Beta vulgaris*) Fields', Iranian Journal of Weed Science, 11(2), pp. 129-143.
- Sarani H, Izadi E, Ghanbari A, Rahemi A. Effect of Temperature and Light on Germination Characteristics of Japanese Morning Glory (*Ipomoea nil*): Determination of Cardinal Temperatures of Germination. Iranian J. Seed Res.. 2019; 6 (1) :115-127
- Abbasi, I., Zaeefarian, F., & Younesabadi, M. (2022). Study of Biological Aspect of Germination and Emergence in Morning Glory (*Ipomoea purpurea* L.). Journal of Iranian Plant Protection Research 36(1): 125-139.
- Dehghan S (2022) Investigating the effect of some environmental factors on seed germination of three new *Ipomoea* species. MSc Thesis, Gorgan University of Agricultural Sciences and Natural Resources, Gorgan, Iran.
- Mina Safavi, Mohammad Rezvani, Faezeh Zaeefarian, Sajede Golmohammadzadeh, and Brian M. Sindel. 2023. Seed germination requirements of *Amaranthus retroflexus* L. populations exposed to environmental factors. Botany. 101(4): 99-111. <https://doi.org/10.1139/cjb-2022-0077>
- Talae, M., Rezvani, M., Radmard, M. & Sindel, B.M. (2024) Influence of environmental factors on seed germination and seedling emergence of *Amaranthus blitoides* S. Watson and *A. hybridus* L. Weed Research, 64(1): 31-41. <https://doi.org/10.1111/wre.1260>
- Talae, M., Rezvani, M., Radmard, M. & Sindel, B.M. (2024) Influence of environmental factors on seed germination and seedling emergence of *Amaranthus blitoides* S. Watson and *A. hybridus* L. Weed Research, 64(1): 31-41. <https://doi.org/10.1111/wre.1260>
- Ghorbani, R. Asgarpour, R. Ghorbani, M. Khajeh, Hosseini, (2014). Effect of environmental factors on wild poinsettia (*Euphorbia heterophylla* L.) germination- a new introduced weed in soybean field, Weed Research Journal, 5(2), 183-198. magiran.com/p1268145
- Ghorbani, R. Asgarpour, R. Ghorbani, M. Khajeh, Hosseini, (2014). Effect of environmental factors on wild poinsettia (*Euphorbia heterophylla* L.) germination- a new introduced weed in soybean field, Weed Research Journal, 5(2), 183-198. magiran.com/p1268145
- Derakhshan, A., Gharekhloo, J. (2014). 'Study on Some Ecological Aspects of Cutleaf groundcherry (*Physalis angulata* L.) Seed Germination and Dormancy', Journal of Iranian Plant Protection Research, 28(3), pp. 416-424. doi: 10.22067/jpp.v28i3.28189
- Sadeghloo, A. & Asghari, J. & Ghaderi-Far, Farshid. (2013). Seed germination and seedling emergence of velvetleaf (*Abutilon theophrasti*) and Barnyardgrass (*Echinochloa crus-galli*). Planta Daninha. 31. 259-266. 10.1590/S0100-83582013000200003.
- Eskandari, B., Ahmadvand, G., Abutalebian, M. A. (2020). 'Effects of salinity and drought stress on germination and seedling *Datura innoxia* Mill.', Environmental Stresses in Crop Sciences, 13(4), pp. 1319-1327. doi: 10.22077/escs.2020.2362.1611
- Sadeqnejad, N., Alizadeh, H., Rahimian Mashhadi, H., Oveisi, M., Tokasi, S. (2023). 'Cardinal temperatures of *Ambrosia* (*Ambrosia psilostachya*) using regression models', Iranian Journal of Weed Science, 19(1), pp. 1-10. doi: 10.22092/ijws.2022.359817.1419
- Mojab, Mehdi & Zamani, Gholam & Eslami, Seyed & Hosseini, Mojtaba & Naseri, S.. (2010). Effects of salt (NaCl) and drought (PEG6000) stresses on germination characteristic and seedling growth of Barnyard grass (*Echinochloa crus-galli* Var: *oryzicola*). 10.22067/jpp.v13i3.24.3856.
-